



MoB

Soluble liquid fertilizer

GET Technology

A SOLUBLE LIQUID FERTILIZER TO SUPPLY NITROGEN, BORON, COPPER, MOLYBDENUM AND ZINC TO CROPS AS LISTED.

14-0-0

Guaranteed analysis:

Nitrogen (N)*	14.00%	Molybdenum (Mo)	0.80%
Boron (B)	6.45%	Zinc (Zn)***	3.35%
Copper (Cu)**	0.15%		

Derived from: Boric acid, Copper(II) hydroxide amino acid complex, Sodium molybdate dihydrate, LB Urea, Zinc oxide amino acid complex.

*Urea Nitrogen - 14.00%

**Chelated Copper - 0.15%

***Chelated Zinc - 3.35%

KEEP OUT OF REACH OF CHILDREN

WARNING: HARMFUL IF SWALLOWED

AVOID INHALATION OR CONTACT WITH EYES OR SKIN

Net Content:

5 gal (19 L)
55 lbs (25 kg)

Density: 11 lbs/gal at 68°F



Manufactured, Distributed and Guaranteed by:
MBFi LLC
4006 S 40th Street, St. Joseph, Missouri, 64503
www.mbf.bio



Nutrition

WARNING / AVISO

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle. (If you do not understand the label, find someone to explain it to you in detail).

This product contains boron; use on any crops other than those recommended may result in serious injury to the crops. Additional supplemental application of boron should not be made unless soil or plant tissue tests indicate that this is necessary.

This fertilizer is to be used only on soil which responds to Molybdenum (Mo). Crops high in Molybdenum (Mo) are toxic to grazing animals (ruminants).

PRECAUTIONARY STATEMENTS

Read the label before use. If medical advice is needed, have a product container or label at hand. Wash hands thoroughly after handling. Wear eye protection and face protection. Avoid breathing fumes/mist/vapors/spray. Use only outdoors or in a well-ventilated area. Wash protective equipment and work clothes thoroughly after handling.

GENERAL INFORMATION

Bio-Finalist MoB is a foliar-applied secondary nutrient. Bio-Finalist MoB should preferably be applied in the early morning or late afternoon. The rate of application is dependent on the reaction required. The lower rates should be used for maintenance while the higher rates will correct nutritional deficiencies.

A pre-mix "jar test" is recommended prior to mixing with any other products to check compatibility.

DIRECTIONS FOR USE

Shake the product thoroughly and dilute it in sufficient water to ensure adequate, even coverage without producing runoff. Agitate the spray mixture during application. Use only as indicated on this label. Clean applicator and mixing containers thoroughly after use and dispose of wash water where it will not contaminate crops, grazing, rivers, dams and boreholes.

IMPORTANT

Because boron is not mobile within the plant, a continuous supply to all growing points is required from germination to maturity. Supplemental application of boron is recommended when soil tests show less than 1.0 ppm boron or tissue test levels fall below average values specified for the crop. Seek advice from a local agricultural authority or the supplier on the recommended application rates.

APPLICATION RATES

Crops	Dosage qt/acre	Remarks
Alfalfa	1	Apply 10 days before cutting.
Bean:		
Dry and Green	1-1.25	Apply at R1-R2 and apply between R7-R8.
Canola	0.25-1	Apply at flowering before pods appear.
Carrot, Garlic, Onion	1-1.75	Apply 3 weeks before harvest.
Corn	1-1.75	Apply at V4-V8 and 15 days before tasselling.
Cotton	1-1.75	Apply 2-3 weeks before defoliation.
Pepper and Tomato	1	Apply 14 days before harvest.
Potato	1-1.75	Apply 2-3 weeks before killing. If vines are not killed, apply 3 weeks before harvest.
Soybean	1-1.25	Apply at stage R1-R2 and at stage R5.1 before pods appear.
Sugar beet	1-1.75	Apply when plants should normally be senescing.
Sunflower	1-1.25	Apply at 20 cm height (knee height) and at head formation.
Wheat	1	Apply 15 days before the seed head appears.

STORAGE AND DISPOSAL

This product must be stored between 40 and 80°F (4 and 27°C). Failure to do so will cause irreversible damage and render the product unfit for use. Use before the expiration date. Store in a cool, dry place out of direct sunlight. Store in the original container only. Keep containers sealed when not in use. Keep the product separate from food and animal forage and out of the reach of children and animals. Do not reuse the empty container. Dispose of rinsate and container in accordance with local, state and federal regulations.

NOTICE OF WARRANTY

CONDITIONS OF SALE-LIMITED WARRANTY AND LIMITATIONS OF LIABILITY AND REMEDIES: To be read prior to using this product. The directions should be followed carefully together with good application practice. Failure to do so may cause injury to crops, animals, man or the environment. Extraordinary or unusual weather conditions may affect product performance and/or crop injury which are beyond the control of MBFI LLC (the "Company") or seller. The Company warrants that this product conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the Directions for Use. The Company makes no other warranties or representations of any kind, express or implied, concerning the product, including no implied warranty of merchantability or fitness for any particular purpose, and no such warranty shall be implied by law. The exclusive remedy against the Company for any cause of action relating to the handling or use of this product shall be limited to, at Company's election, ONE of the following: 1. Refund of the purchase price paid by buyer or user for product bought. 2. Replacement of the product used. To the extent allowed by law, the Company shall not be liable and any and all claims against the Company are waived for special, indirect, incidental, or consequential damages or expense of any nature, including, but not limited to, loss of products or income. The company/seller offer this product and the buyer/user accept it, subject to the foregoing conditions of sale and limitation of warranty, liability and remedies.